

TRANSMISSION OF INFORMATION BY EXTRATERRESTRIAL CIVILIZATIONS

N. S. Kardashev

P. K. Shternberg Astronomical Institute

Translated from *Astronomicheskii Zhurnal*, Vol. 41, No. 2,

pp. 282-287, March-April, 1964

Original article submitted December 12, 1963

The protracted duration of signal propagation is a determining factor in the one-way transmission of information through space. Reliable reception, or any reception at all, of signals by unknown subscribers necessarily requires an isotropic emission. The optimum signal spectrum for transmitting the maximum amount of information in the presence of quantum noise and the background of cosmic radio-frequency emission has been calculated. It is shown that a civilization located at any distance in the universe and in possession of power on the order of $L_{\odot} \approx 4 \times 10^{33}$ erg/sec or more, which it is capable of transmitting in a coded isotropic radio-frequency signal, may be detected by conventional radio astronomical techniques. The expected distinguishing properties of artificial sources of cosmic radio-frequency emission are enumerated. It is speculated that even some sources known to us today (notably CTA-21 and CTA-102) may be artificial radio sources.

1. The principal factors which exert a determining effect on the range of space radio communications are the transparency of the interstellar medium to radio signals, the level of the equipment noise and space noise, and the power of the transmitters. The greatest possible range for establishing space communications could be set most likely in the range from 10^9 to 10^{11} cps [1]. The absorption coefficient of the interstellar medium is negligibly small at those frequencies. The equivalent noise temperature may be represented in the form $T_N = T_n + T_t + T_q$, where T_n and T_t are respectively the temperature due to synchrotron radiation and due to background thermal cosmic radio emission, and $T_q = hf/k$ is the equivalent noise temperature due to quantum fluctuations in the minimum detectable signal (h and k are the Planck constant and the Boltzmann constant). The expression for T_N gives an estimate of the limiting sensitivity which might be achieved in the case of an ideal noise-free receiver and observations outside the earth's atmosphere. In Fig. 1, we find plots of T_N as a function of the frequency in accord with up-to-date radio astronomy data [2]. The paramount role in the establishing of long-range communications within the confines of our galaxy will evidently be played by thermal and nonthermal radio-frequency emission from the

galactic disk (over a range of $\pm 50^\circ$ in longitude on either side of the center of the galaxy). In that case, we have

$$T_N = 2 \cdot 10^{27} \cdot f^{-2.9} + 10^{19} f^{-2} + 4.8 \cdot 10^{-14} f. \quad (1)$$

In dealing with the problem of possible success in setting up communications between the galaxies, we must take into consideration the brightness tem-




Fig. 1. Noise spectrum outside the confines of the earth's atmosphere: —) In the direction pointing toward the center of the Galaxy; ----) in the direction toward the galactic pole.

perature of the background at high galactic latitudes, which is due to synchrotron radiation from the halo and from the metagalaxy. In this case, we have

$$T_N = 10^{26} f^{-2.9} + 4.8 \cdot 10^{-11} f. \quad (2)$$

In both cases, the noise temperature will display a deep-sloping minimum in the decimeter and centimeter wavelength ranges, which renders this range more suitable for space communications over exceptionally vast distances.

2. Let us evaluate the information content of communications channels for application to this problem.

The upper bound of the rate of information transmission at a specified average transmitter power and specified noise distribution is determined by the corresponding Shannon theorem [3]:

$$R = \int_{f_1}^{f_2} \log_2 \left[\frac{S(f) + n(f)}{n(f)} \right] df, \quad (3)$$

where $S(f)$, $n(f)$ are the functions of the spectral power density of the useful signal and of the noise, respectively. By solving the appropriate variational problem, we may show the maximum rate of information transmission to be achieved under the condition

$$S(f) + n(f) = n(f_1) = n(f_2). \quad (4)$$

Here, f_1 and f_2 are the bounds of the transmitter transmission band. It is accordingly quite clear that the spectrum of the artificial source must display the shape of the curve in Fig. 1, but with the sign reversed,

$$S(f) = n(f_1) - n(f),$$

i. e., the spectrum of the artificial radio emission must feature a maximum, and in a region of frequencies lower than this frequency will fall as $a - bf^{-2.9}$, whereas, in the region of higher frequencies, it will fall as $a - cf$ (a , c , b here are constants dependent upon the power and bandwidth of the transmitters and on the noise distribution).

To make a rough estimate of the rate of information transmission, there is of course no need to take the logarithmic term in Shannon's theorem into account. The rate of information transmission $R = f_2 - f_1 = \Delta f$ then, i. e., it will be equal to the system bandwidth. Let P be the transmitter power, and we shall assume the radiation isotropic. There appears to be no point, in our view, in discussing the possibility of establishing outer space communications when high-directivity antennas [4] are

being used for transmission, since the probability of success in establishing such communications is virtually nil. The high directivity of the radiation can be conveniently used, in all probability, only after a two-way communication exchange has been set up, and if we bear in mind the fact that the distances between the two civilizations in question may be comparable to or even more enormous than the dimensions of our galaxy, then this second mode of communications may be brought to fruition much later, and in the meantime the need for isotropic radiation geared to establish contacts with new potential listeners will not have abated. Let the transmission be carried out over a bandwidth Δf , and let the maximum expected distance of the transmission be r , with A the effective area presented by the receiving antenna, T_N the noise temperature referred to the input of the receiving antenna. We shall assume the transmission to be reliable when the condition $100 kT_N = PA/4\pi r^2 \Delta f$ is fulfilled, i. e., when there is a 100-fold excess of signal over noise. Then

$$\Delta f = PA / 400\pi r^2 kT_N. \quad (5)$$

Consequently, the channel capacity will not be less than Δf bits per second. The radio emission flux per unit frequency interval will not be less than

$$F_f = 100 kT_N / A. \quad (6)$$

3. The most important parameter to deal with in this problem is the power P , about which several hypotheses are in order. Calculations show that the total quantity of energy expended by all of mankind per second at the present time is about 4×10^{19} erg, and the annual increase in this energy expenditure is placed at 3-4% over the next 60 years, on the basis of statistical findings [5]. Now let this increment represent an annual increase in energy consumption by a factor of $1 + x$, so that in t years the increase will be $(1 + x)^t \approx e^{tx}$ times ($x \ll 1$). Assuming $x = 1\%$, we find that the energy consumption per second will be equal to the output of the sun per second, 3200 years from now, i. e., 4×10^{33} erg/sec, and that in 5800 years the energy consumed will equal the output of 10^{11} stars like the sun. The figures arrived at seem to be inordinately high when compared to the present level of development, but we see no reasons why the tempo of increase in energy consumption should fall substantially than predicted. Moreover, the availability of a large amount of information forthcoming from other and more highly developed civilizations might

contribute to a staggering increase in energy consumption.

In line with the estimates arrived at, it will prove convenient to classify technologically developed civilizations in three types:

I – technological level close to the level presently attained on the earth, with energy consumption at $\approx 4 \times 10^{19}$ erg/sec.

II – a civilization capable of harnessing the energy radiated by its own star (for example, the stage of successful construction of a "Dyson sphere" [6]); energy consumption at $\approx 4 \times 10^{33}$ erg/sec.

III – a civilization in possession of energy on the scale of its own galaxy, with energy consumption at $\approx 4 \times 10^{44}$ erg/sec.

4. Estimates of the possibility of detecting a type I civilization [7] and related experiments in the "OZMA" project in the USA have revealed the extremely low probability of any such event. Consider the possibility of detection and reception of information sent by type II and type III civilizations. First of all, we assume here that one of the principal tasks of such communication efforts would be the transmission of information from a more highly developed civilization to a less highly developed one. Starting from the present level of development of radio physics as point of departure, we see that in principle it is possible to build antennas, within the next two decades, with an effective area of 10^5 m² and with receiving apparatus featuring a noise temperature $T_N \approx 1^\circ\text{K}$. If a transmitter is designed for this system to receive and record information, Eq. (3) will show that the radio emission flux at the receiving point will be not less than 1.4×10^{-26} W/m² · cps, an amount which is well within the recording capabilities of presently existing radio telescopes. [Of course, far simpler equipment will be required to detect the signals since, in contrast to the reception of information, here we may utilize the averaging techniques common in radio astronomy, and the radiometric gain $\sqrt{\Delta f \tau}$ (where τ is the build-up time) may enhance the sensitivity by

3 to 4 orders of magnitude over a fairly wide signal band.]

We now cite some estimates of the quantity of information obtainable, in line with formula (2). For a type II civilization, we have to take into account the frequency variation of T_N as in accord with formula (1). $T_N = 1^\circ\text{K}$ is assumed for a type II civilization. The table lists estimates for three distances corresponding to the transmission of information within the confines of the galaxy, within the confines of a local system of galaxies, and within the confines of the portion of the metagalaxy accessible to observation. The estimates arrived at show that should there exist even one type II civilization within the confines of the local system of galaxies, there will be a realistic possibility of securing an enormous quantity of information. The same holds for the existence of even one single type III civilization in the portion of the universe accessible to observation. For purposes of comparison, let us estimate the time required to transmit 10^8 printed and manuscript-form publications now available on the earth through a channel of 10^9 cps bandwidth. Assuming that each written work contains an average of about 10^6 bits of information, we find that the total quantity of 10^{14} bits of information may be transmitted in 10^5 sec, i.e., in a single day. However, it is quite evident that there is no need to transmit all of the 100 million publications in order to broadcast the principal data on the status of science, technology, and culture on the earth, for such information would contain a colossal amount of "redundancy." Apparently, all of the basic information could be compressed in 10^5 books of 10^6 bits each, which would come to 10^{11} bits and would take only 100 sec to be transmitted via the same communications channel. Finally, it is entirely reasonable to assume that type II and type III civilizations would be in possession of information many orders of magnitude in excess of what we have available at the present time. For that reason, they would have to be broadcasting practically continual-

Number of Bits per Second $\sim \Delta f$

Type of civilization	Transmitter power	$r = 100,000$ light years	$r = 10$ million light years	$r = 10$ billion light years
II	4×10^{33} erg/sec	3×10^9	3×10^5	Transmission of a large quantity of information impossible
III	4×10^{44} erg/sec	2.4×10^{15}	2.4×10^{13}	$3 \cdot 10^{10}$

ly, and this would also be the case for increasing the possibility of reception by type I civilizations. Moreover, in order to improve the reliability of the information received and in order to afford some opportunity to make connections with new subscribers, there would have to be periodic repetitions of the programs broadcast.

Note again that, in all likelihood, a type I civilization would be capable of sending a return signal only after its energy consumption had increased measurably. Consequently, the communications would be a one-way affair at the start, and the problem of how long it takes the signals to propagate would be a secondary one.

As is evident from the above estimates of transmitter power for type II and type III civilizations, the figures are very close to the power of synchrotron radiation from nebulas formed in supernova explosions, or from radio galaxies. Calculations of the optimum transmitter bandwidth show that the transmission spectrum may also closely resemble the spectra of discrete radio sources. Several criteria could be singled out which would be useful in discriminating artificial radio sources from the vast number of radio stars accessible to observation.

The artificial sources would evidently 1) have to have very small angular dimensions (at least in the case of type II civilizations); the angular dimensions would have to be of the order of the angular dimensions of stars, i. e., less than $0''.001$. The now known natural radio sources must have appreciably larger angular dimensions, larger than $0''.01$ in fact, according to theory [8]; 2) they would have to possess circular polarization, so that the effect of the Faraday rotation of the plane of polarization in the interstellar medium would not distort the information received; 3) they would have to exhibit variability in time without leading to statistical fluctuations; this is obviously a criterion of outstanding importance, but it is possible that a regularity in time variations of the signal might be revealed only in the course of observations using special equipment of wide bandwidth (10^9 cps) and of sufficient sensitivity to operate with very short time constants; 4) finally, it is to be anticipated that certain details would be present in the spectrum of the source suspected of artificiality which would have been designed for the express purpose of emphasizing its artificial origin; in particular, we might anticipate such a feature in the environs of the 21-cm wavelength line. Information transmission at that frequency within the confines of this galaxy would be inadvisable, since the signal will be strongly

absorbed by neutral hydrogen. It would be reasonable, for that reason, to eliminate a band of 1-2 Mc width, say of rectangular shape, in the continuous spectrum of the artificial source, to lay special emphasis on the unusual nature of the radiation. Considerations involving the anticipated shape of the complete spectrum of the artificial source also deserve close attention. These arguments were cited above when the estimate was made of the quantity of information which could be transmitted. The most characteristic feature of the spectrum is the linear dependence of the flux on frequency in the high-frequency region of the spectrum (cf. Fig. 2).

It is consequently of the utmost importance to carry out a program of studying and searching discrete radio sources, in the immediate future, with due attention paid to the spectral features mentioned here. Note that even at the present time we have knowledge of about twenty to thirty or so radio sources, with the upper limit of the angular dimensions in the range of 1 to 10 sec of arc [9]. Some of these sources have been identified with peculiar optical objects the nature of which is still obscure [10]. Most of them have not yet been successfully identified with optical objects.

For example, two sources of radio-frequency emission from outer space, CTA-21 and CTA-102, were recently discovered at the California Technological Institute [11], and display angular dimensions not less than $20''$, and have not been identified with a single one of the optical objects in the Palomar




Fig. 2. ———) Anticipated emission spectrum of radio transmitters of extraterrestrial civilizations; - - -) spectrum of radio sources CTA-21 and CTA-102, suspected of being artificial radio sources, and spectrum of a typical natural radio source Virgo A.

sky charts and, even more intriguing, these sources exhibit a spectrum highly similar to the anticipated artificial spectrum. Figure 2 presents the observational data reported in [12] on these two sources, alongside the theoretically predicted spectrum. For purposes of comparison, the spectrum of a typical natural radio source, Virgo A, is added (the scale is compressed by a factor of 10 on the ordinate axis in the case of Virgo A).

The most promising region for a search for artificial radio signals is apparently that in the direction toward the center of the Galaxy, since the density of the stellar population is greatest there along the line of sight. It would also be appropriate to investigate the closest galaxies, and in the first instance the large nebula in the Andromeda constellation and the Magellanic Clouds, as well as the closest radio galaxies NGC 4486 and NGC 5128.

In conclusion, we should like to note that the estimates arrived at here are unquestionably of no more than a tentative nature. But all of them bear witness to the fact that, if terrestrial civilization is not a unique phenomenon in the entire universe, then the possibility of establishing contacts with other civilizations by means of present-day radio physics capabilities is entirely realistic. At the same time, it is very difficult to accept the notion that, of all of the 10^{11} stars present in our Galaxy, only near the sun has a civilization developed. It is still more difficult to extend this inference to the 10^{10} galaxies existing in the portion of the universe accessible to observation. In any case, the deciding word on this question is left to experimental verification. In particular, we may anticipate that space rockets will clear up the question of whether or not life exists on other planets in the solar system in the years to come. The discovery of even the very simplest organisms, on Mars for instance, would greatly increase the probability that many type II civiliza-

tions exist in the Galaxy. Radio astronomical searches could of course play a decisive part in resolving this problem.

LITERATURE CITED

1. B. M. Oliver, *International Science and Technology*, No. 10, 55, October, 1962.
2. A. J. Turtle, J. F. Pugh, S. Kenderdine, and J. J. K. Pauliny-Toth, *Monthly Notices Roy. Astron. Soc.*, 124, 297 (1962); R. W. Wilson, *Observations of the Owen Valley Radio Observ.*, Cal. Techn., No. 3 (1963); W. Altenhoff, P. G. Mezger, H. Strasse, H. Wendkerr, and G. Westerhout, *Ver entliche Univ. - Sternvarte zu Bohn*, No. 9 (1960).
3. S. Goldman, *Information Theory*, Prentice-Hall, New York (1953).
4. G. Cocconi and Ph. Morrison, *Nature*, 184, 844 (1959).
5. P. C. Putnam, *Energy in the Future*, New York (1948).
6. F. G. Dyson, *Science*, 131, 1667 (1959).
7. F. D. Drake, *Sky and Telescope*, 19, 140 (1959).
8. V. I. Slysh, *Nature*, 199, 682 (1963).
9. R. L. Allen, B. Anderson, R. G. Conway, H. P. Palmer, V. C. Reddich, and B. Rowson, *Monthly Notices Roy. Astron. Soc.*, 124, 477 (1962).
10. M. Schmidt, *Nature*, 197, 1040 (1963); J. B. Oke, *Nature*, 197, 1042 (1963); J. Greenstein and T. A. Matthews, *Nature*, 197, 1042 (1963).
11. D. E. Harris and J. A. Roberts, *Publ. Astron. Soc. Pacific* 72, 237 (1960).
12. R. G. Conway, K. I. Kellerman, and R. J. Long, *Monthly Notices Roy. Astron. Soc.*, 125, 261 (1963).